THE CANADIAN ENTOMOLOGIST

Index to Volume 128 (1996)

SUPPLEMENTS AND MEMOIRS

THE ENTOMOLOGICAL SOCIETY OF CANADA, OTTAWA

Albages, R. See Buès et al., 849.
AliNiazee, M.T. See Kasana and AliNiazee, 377.
Baoua Boukary, I., J.-C. Tourneur, and J. Gingras. Life cycle of *Forficula senegalensis* Serv. (Dermaptera: Forficulidae) and its relationship to the development of bulrush millet in the Sudanese-Sahelian zone of Niger, 831.
Bauce, E. See Han and Bauce, 167.
Berryman, A.A. See Valenti et al., 839.
Blatt, S.E., and J.H. Borden. Distribution and impact of *Leptoglossus occidentalis* Heidemann (Hemiptera: Coreidae) in seed orchards in British Columbia, 1065.
Evidence for a male-produced aggregation pheromone in the western conifer seed bug, *Leptoglossus occidentalis* Heidemann (Hemiptera: Coreidae), 777.
Bodnaryk, R.P. See Xie et al., 865.
Boiteau, G., and W. Coleman. Cold tolerance in the Colorado potato beetle, *Leptinotarsa decemlineata* (Say) (Coleoptera: Chrysomelidae), 1087.
Borden, J.H. See Ntiamoah and Borden, 351.
See Tomlin and Borden, 539.
See Blatt and Borden, 777.
See McBrien et al., 1057.
See Blatt and Borden, 1065.
Bousquet, Y. Taxonomic revision of Nearctic, Mexican, and West Indian Oodini (Coleoptera: Carabidae), 443.
Brandt, J.P., S.M. Smith, and M. Hubbes. Distribution and sampling of root weevil larvae in young ornamental conifer plantations, 1125.
Brennan, P.A. See Shishehbor and Brennan, 859.
Broatch, J.S. See Vernon and Broatch, 1077.
Bros, W.E. See Hoover et al., 981.
Buès, R., M. Eizaguirre, J.F. Toubon, and R. Albages. Différences enzymatiques et écophysiologiques entre populations de *Sesamia nonagrioïdes* Lefèbre (Lépidoptère: Noctuidae) originaires de l'ouest du bassin méditérranéen, 849.
Burban, C. See Jactel et al., 1143.
Butler, L. See Sample et al., 573.
Chakrabarti, S., and F.W. Quednau. A new genus, *Neothelaxes*, and two new species of Thelaxinae from India (Homoptera: Aphididae), 1005.
Çilgi, T., S.D. Wratten, J.L. Robertson, D.E. Turner, J.M. Holland, and G.K. Frampton. Residual toxicities of three insecticides to four species (Coleoptera: Carabidae) of arthropod predators, 1115.
Coleman, W. See Boiteau and Coleman, 1087.
Cowle, N.T. See Dosdall et al., 1157.
Currie, C.R., J.R. Spence, and W.J.A. Volney. Biology and life history of *Epuraea obliquus* Hatch (Coleoptera: Nitidulidae) on western gall rust, 177.
Denis, J. See Masner and Denis, 85.
Dosdall, L.M., M.J. Herbut, N.T. Cowle, and T.M. Micklich. The effect of tillage regime on emergence of root maggots (*Delia* spp.) (Diptera: Anthomyiidae) from canola, 1157.
Eizaguirre, M. See Buès et al., 849.
Elliott, R.H., and L.W. Mann. Susceptibility of red spring wheat, *Triticum aestivum* L. cv. Katepwa, during heading and anthesis to damage by wheat midge, *Sitodiplosis mosellana* (Géhin) (Diptera: Cecidomyiidae), 367.
Eveleigh, E.S. See Piene and Eveleigh, 1101.
See Huber et al., 1167.
Feng, R. See Isman et al., 353.
Ferrell, G.T. See Valenti et al., 839.
Fidgen, L.L., and J.D. Sweeney. Fir coneworm, *Dioryctria abietivorella* (Groté) (Lepidoptera: Pyralidae), prefer cones previously exploited by the spruce cone maggots *Strobilomyia neanthracina* Michelsen and *Strobilomyia appalachensis* Michelsen (Diptera: Anthomyiidae), 1221.
Fields, P.G. See Xie et al., 865.
Fitzpatrick, S.M. See Li and Fitzpatrick, 605.
Fox, J.W. See Hoover et al., 981.
Frampton, G.K. See Çilgi et al., 1115.
Gilbert, N., and D.A. Raworth. FORUM: Insects and temperature—a general theory, 1.
Gingras, J. See Baoua Boukary et al., 831.
Gray, T.G., R.F. Shepherd, G. Gries, and R. Gries. Sex pheromone component of the western blackheaded budworm, *Acleris gloverana* Walsingham (Lepidoptera: Tortricidae), 1135.
Greco, C.F., D. Holland, and P.G. Kevan. Foraging behaviour of honey bees (*Apis mellifera* L.) on staghorn sumac [*Rhus hirta* Sudworth (ex-*typhina* L.)]: Differences and dioecy, 355.
Gries, G. See Gray et al., 1135.

Gries, R. See Gray et al., 1135.

Guo, J.-Q., J.-P. Moreau, and H. Lapierre. Variability among aphid clones of *Rhopalosiphum padi* L. and *Sitobion avenae* Fabr. (Homoptera: Aphididae) in transmission of three PAV isolates of barley yellow dwarf viruses, 209.

Han, E.-N., and E. Bauce. Diapause development of spruce budworm larvae, *Choristoneura fumiferana* (Clem.) (Lepidoptera: Tortricidae), at temperatures favouring post-diapause development, 167.

Harris, P., and J.D. Shorthouse. Effectiveness of gall inducers in weed biological control, 1021.

Harvey, G.T. Population genetics of the spruce budworm, *Choristoneura fumiferana* (Clem.) Freeman (Lepidoptera: Tortricidae), in relation to geographical and population density differences, 219.

Genetic relationships among *Choristoneura* species (Lepidoptera: Tortricidae) in North America as revealed by isozyme studies, 245.

Heinz, K.M. Space- and cohort-dependent longevity in adult *Liriomyza trifolii* (Burgess) (Diptera: Agromyzidae) mass-rearing cultures, 1225.

Herbut, M.J. See Dosdall et al., 1157.

Holland, D. See Greco et al., 355.

Holland, J.M. See Çilgi et al., 1115.

Holliday, N.J. See Pernal et al., 791.

Hoover, K., D.L. Wood, A.J. Storer, J.W. Fox, and W.E. Bros. Transmission of the pitch canker fungus, *Fusarium subglutinans* f. sp. *pini*, to Monterey pine, *Pinus radiata*, by cone- and twig-infesting beetles, 981.

Horton, D.R., T.M. Lewis, and L.G. Neven. Reduced cold-hardiness of pear psylla (Homoptera: Psyllidae) caused by exposure to external water and surfactants, 825.

Howden, A.T. Neotropical *Pandeleteius* (Coleoptera: Curculionidae) with irregular elytral striae, 877.

Hubbes, M. See Brandt et al., 1125.

Huber, J.T. See Ikeda and Huber, 743.

See Ikeda et al., 767.

Huber, J.T., E. Eveleigh, S. Pollock, and P. McCarthy. The chalcidoid parasitoids and hyperparasitoids (Hymenoptera: Chalcidoidea) of *Choristoneura* species (Lepidoptera: Tortricidae) in America north of Mexico, 1167.

Hunt, D.W.A., and G. Whitfield. Potato trap crops for control of Colorado potato beetle (Coleoptera: Chrysomelidae) in tomatoes, 407.

Ikeda, E., and J.T. Huber. Review of the world species of *Dimmockia* Ashmead (Hymenoptera: Eulophidae), 743.

Ikeda, E., K. Kamijo, and J.T. Huber. A new genus of Tetrastichinae (Hymenoptera: Eulophidae) from Japan, 767.

Ishiwata, S.-I. A study of the genus *Ephoron* from Japan (Ephemeroptera, Polymitarcyidae), 551.

Isman, M.B., R. Feng, and D.L. Johnson. Detoxicative enzyme activities in five species of field-collected melanopline grasshoppers (Orthoptera: Acrididae), 353.

Ivie, M.A., and A.G. Zinovjev. Discovery of the wheat stem sawfly (*Cephus cinctus* Norton) (Hymenoptera: Cephidae) in Asia, with the proposal of a new synonymy, 347.

Jactel, H., N. Perthuisot, P. Menassieu, G. Raise, and C. Burban. A sampling design for within-tree larval populations of the maritime pine scale, *Matsucoccus feytaudi* Duc. (Homoptera: Margarodidae), and the relationship between larval population estimates and male catch in pheromone traps, 1143.

Johnson, D.L. See Isman et al., 353.

Johnson, N.F. Revision of world species of *Paratelenomus* Dodd (Hymenoptera: Scelionidae), 273.

Judd, G.J.R. See McBrien et al., 1057.

Kamijo, K. See Ikeda et al., 767.

Kasana, A., and M.T. AliNiazee. Seasonal phenology of the walnut husk fly, *Rhagoletis completa* Cresson (Diptera: Tephritidae), 377.

Kevan, P.G. See Holland et al., 355.

Lapierre, H. See Guo et al., 209.

Larson, D.J. Revision of North American *Agabus* Leach (Coleoptera: Dytiscide): The *opacus*-group, 613.

Lemmon, C.R. See Maier and Lemmon, 995.

Le Ralec, A., J.M. Rabasse, and E. Wajnberg. Comparative morphology of the ovipositor of some parasitic Hymenoptera in relation to characteristics of their hosts, 413.

LeSage, L., and P. Paquin. Identification keys for *Aphthona* flea beetles (Coleoptera: Chrysomelidae) introduced in Canada for the control of spurge (*Euphorbia* spp., Euphorbiaceae), 593.

Lewis, T.M. See Horton et al., 825.

Li, S.Y., and S.M. Fitzpatrick. The effects of application rate and spray volume on efficacy of two formulations of *Bacillus thuringiensis* Berliner var. *kurstaki* against *Choristoneura rosaceana* (Harris) (Lepidoptera: Tortricidae) on raspberries, 605.

Linton, D.A. See Safranyik et al., 199.

Lyons, D.B. Oviposition and fecundity of pine false webworm (Hymenoptera: Pamphiliidae), 779.

Mackauer, M. See Stadler and Mackauer, 27.

Mackauer, M., J.P. Michaud, and W. Völkl. INVITATION PAPER: C.P. Alexander Fund. Host choice by aphidiid parasitoids (Hymenoptera: Aphidiidae): Host recognition, host quality, and host value, 959.
Maier, C.T., and C.R. Lemmon. Life cycle of *Lambdina athasaria* (Walker) (Lepidoptera: Geometridae) developing on hemlock in Connecticut, 995.
Mann, L.W. See Elliott and Mann, 367.
Masner, L., and J. Denis. The Nearctic species of *Idris* Foerster. Part I: The *melleus*-group (Hymenoptera: Scelionidae), 85.
McBrien, H.L., G.J.R. Judd, and J.H. Borden. Potential for pheromone-based mating disruption of the mullein bug, *Campylomma verbasci* (Meyer) (Heteroptera: Miridae), 1057.
McCarthy, P. See Huber et al., 1167.
McClay, A.S. Unisexual broods in the gall midge *Cystiphora sonchii* (Bremi) (Diptera: Cecidomyiidae), 775.
McClure, M., D.T. Quiring, and J.J. Turgeon. Oviposition, temporal distribution, and potential impact of *Strobilomyia laricis* Michelsen and *S. viaria* (Huckett) (Diptera: Anthomyiidae) on eastern larch, *Larix laricina* (Du Roi) K. Koch, 67.
Melathopoulos, A.P., M.L. Winston, J.S. Pettis, and T. Pankiw. Effect of queen mandibular pheromone on initiation and maintenance of queen cells in the honey bee (*Apis mellifera* L.), 263.
Menassieu, P. See Jactel et al., 1143.
Michaud, J.P. See Mackauer et al., 959.
Micklich, T.M. See Dosdall et al., 1157.
Milbrath, L.R. See Nead et al., 349.
Moreau, J.-P. See Guo et al., 209.
Nead, B.A., M.J. Weiss, and L.R. Milbrath. Feeding preferences of the ashgray blister beetle, *Epicauta fabricii* (LeConte) (Coleoptera: Meloidae), for four legumes, 349.
Nealis, V.G., D. Oliver, and D. Tchir. The diapause response to photoperiod in Ontario populations of *Cotesia melanoscela* (Ratzeburg) (Hymenoptera: Braconidae), 41.
Neven, L.G. See Horton et al., 825.
Ntiamoah, Y.A., and J.H. Borden. Monoterpene oviposition deterrents for cabbage maggots, *Delia radicum* (L.) (Diptera: Anthomyiidae), 351.
O'Hara, J.E. Earwig parasitoids of the genus *Triarthria* Stephens (Diptera: Tachinidae) in the New World, 15.
The tachinid taxa of Louis P. Mesnil, with notes on nomenclature (Insecta: Diptera), 115.
Oliver, D. See Nealis et al., 41.
Palacios-Vargas, J.G. New species of *Palmanura* (Collembola: Neanuridae) from Mexico and Guatemala, 805.
Pankiw, T. See Melathopoulos et al., 263.
Paquin, P. See LeSage and Paquin, 593.
Peck, S.B., and K. Stephan. A revision of the genus *Colon* Herbst (Coleoptera; Leiodidae; Coloninae) of North America, 667.
Pernal, S.F., D.G. Senanayake, and N.J. Holliday. Patterns of feeding injury to potato by the potato flea beetle (Coleoptera: Chrysomelidae) in Manitoba, 791.
Perthuisot, N. See Jactel et al., 1143.
Pettis, J.S. See Melathopoulos et al., 263.
Philippe, G. See Roques et al., 391.
Piene, H. Changes in spruce budworm defoliation with crown level, 1109.
Piene, H., and E.S. Eveleigh. Spruce budworm defoliation in young balsam fir: The 'green' tree phenomenon, 1101.
Pines, I.L., and A.R. Westwood. Evaluation of monosodium methane arsenate for the suppression of native elm bark beetles, *Hylurgopinus rufipes* (Eichhoff) (Coleoptera: Scolytidae), 435.
Polavarapu, S., and W.D. Seabrook. Seasonal diapause development, effects of temperature and photoperiod on postdiapause egg development, and validation of a degree-day model predicting larval eclosion of blueberry leaftier, *Croesia curvalana* (Kearfott) (Lepidoptera: Tortricidae), 187.
Pollock, S. See Huber et al., 1167.
Quednau, F.W. See Chakrabarti and Quednau, 1005.
Two new species of myzocallidine aphids from North America and Japan (Homoptera: Aphididae), 1013.
Quiring, D.T. See McClure et al., 67.
Rabasse, J.M. See Le Ralec et al., 413.
Raimbault, J.-P. See Roques et al., 391.
Raise, G. See Jactel et al., 1143.
Raworth, D.A. See Gilbert and Raworth, 1.
Reardon, R. See Sample et al., 573.
Richards, K.W. Effect of environment and equipment on productivity of alfalfa leafcutter bees (Hymenoptera: Megachilidae) in southern Alberta, Canada, 47.
Richards, P.C., and J.M. Schmidt. The effects of selected dietary supplements on survival and reproduction of *Orius insidiosus* (Say) (Hemiptera: Anthocoridae), 171.
Robertson, J.L. See Çilgi et al., 1115.

Roques, A., J.-H. Sun, X.-D. Zhang, G. Philippe, and J.-P. Raimbault. Effectiveness of trunk-implanted acephate for the protection of cones and seeds from insect damage in France and China, 391.
Safranyik, L., T.L. Shore, and D.A. Linton. Ipsdienol and lanierone increase *Ips pini* Say (Coleoptera: Scolytidae) attack and brood density in lodgepole pine infested by mountain pine beetle, 199.
Sample, B.E., L. Butler, C. Zivkovich, R.C. Whitmore, and R. Reardon. Effects of *Bacillus thuringiensis* Berliner var. *kurstaki* and defoliation by the gypsy moth [*Lymantria dispar* (L.) (Lepidoptera: Lymantriidae)] on native arthropods in West Virginia, 573.
Sanders, C.J. Effects of prolonged exposure to different concentrations of synthetic pheromone on mating disruption of spruce budworm moths in a wind tunnel, 57.
Sandoval, C.P., and V.R. Vickery. *Timema douglasi* (Phasmatoptera: Timematodea), a new parthenogenetic species from southwestern Oregon and northern California, with notes on other species, 79.
Schmidt, J.M. See Richards and Schmidt, 171.
Seabrook, W.D. See Polavarapu and Seabrook, 187.
Senanayake, D.G. See Pernal et al., 791.
Shepherd, R.F. See Gray et al., 1135.
Shishehbor, P., and P.A. Brennan. Adult longevity, fecundity, and population growth rates for *Trialeurodes ricini* Misra (Homoptera: Aleyrodidae) at different constant temperatures, 859.
Shore, T.L. See Safranyik et al., 199.
Shorthouse, J.D. See Harris and Shorthouse, 1021.
Smith, S.M. See Brandt et al., 1125.
Spence, J.R. See Currie et al., 177.
Stadler, B., and M. Mackauer. Influence of plant quality on interactions between the aphid parasitoid *Ephedrus californicus* Baker (Hymenoptera: Aphidiiidae) and its host, *Acyrthosiphon pisum* (Harris) (Homoptera: Aphididae), 27.
Storer, A.J. See Hoover et al., 981.
Sun, J.-H. See Roques et al., 391.
Sweeney, J.D. See Fidgen and Sweeney, 1221.
Tchir, D. See Nealis et al., 41.
Tomlin, E.S., and J.H. Borden. Feeding responses of the white pine weevil, *Pissodes strobi* (Peck) (Coleoptera: Curculionidae), in relation to host resistance in British Columbia, 539.
Toubon, J.F. See Buès et al., 849.
Tourneur, J.-C. See Baoua Boukary et al., 831.
Turgeon, J.J. See McClure et al., 67.
Turner, D.E. See Çilgi et al., 1115.
Valenti, M.A., A.A. Berryman, and G.T. Ferrell. Arthropods associated with a manzanita gall induced by the aphid *Tamalia coweni* (Cockerell) (Homoptera: Aphididae), 839.
Vernon, R.S., and J.S. Broatch. Responsiveness of *Delia* spp. (Diptera: Anthomyiidae) to colored sticky traps in flowering and rosette stage canola, 1077.
Vickery, V.R. See Sandoval and Vickery, 79.
Völkl, W. See Mackauer et al., 959.
Volney, W.J.A. See Currie et al., 177.
Wajnberg, E. See Le Ralec et al., 413.
Weiss, M.J. See Nead et al., 349.
Westwood, A.R. See Pines and Westwood, 435.
Whidden, T.L. The fidelity of commercially reared colonies of *Bombus impatiens* Cresson (Hymenoptera: Apidae) to lowbush blueberry in southern New Brunswick, 957.
Whitfield, G. See Hunt and Whitfield, 407.
Whitmore, R.C. See Sample et al., 573.
Winston, M.L. See Melathopoulos et al., 263.
Wood, D.L. See Hoover et al., 981.
Wratten, S.D. See Çilgi et al., 1115.
Xie, Y.S., R.P. Bodnaryk, and P.G. Fields. A rapid and simple flour-disk bioassay for testing substances active against stored-product insects, 865.
Zhang, X.-D. See Roques et al., 391.
Zinovjev, A.G. See Ivie and Zinovjev, 347.
Zivkovich, C. See Sample et al., 573.
Zloty, J. A revision of the Nearctic *Ameletus* mayflies based on adult males, with descriptions of seven new species (Ephemeroptera: Ameletidae), 293.